

New perspectives on geproci sets

Jake Kettinger

Colorado State University

20 June 2024

What is Geproci?

Definition

A finite set Z in \mathbb{P}_k^n is **geproci** if the projection \bar{Z} of Z from a general point P to a hyperplane $H = \mathbb{P}_k^{n-1}$ is a complete intersection in H .


Geproci stands for **general projection is a complete intersection**. The only nontrivial examples known are for $n = 3$. In this case a hyperplane is a plane. A reduced set of points in a plane is a complete intersection if it is the transverse intersection of two algebraic curves, [like this](#).

For $\#Z = ab$ ($a \leq b$), Z is (a, b) -geproci if \bar{Z} is the intersection of a degree a curve and a degree b curve.

Trivial Cases: Coplanar Points and Grids

A set of coplanar points in \mathbb{P}^3 is geproci if and only if it is already a complete intersection in the plane containing it.

The easiest non-coplanar examples are grids, which are sets of points that form the intersection of two families of mutually-skew lines.




Summary of Nontrivial Cases

Half-Grids: A procedure is known for creating an (a, b) -geproci half-grid for $4 \leq a \leq b$, but it is not known what other examples there can be.

Non-Half-Grids: Until recently, only a few examples were known and there was no known way to generate more.



Because of this, nontrivial non-half-grids have been mysterious; it's easier to get an idea of what a half-grid is like.



The D_4 configuration is a $(3, 4)$ -geproci half-grid.

Cones and Geproci

It is interesting when there is a cone through Z whose vertex is a general point P , and which meets H in a curve containing the projected image of Z . For Z to be (a, b) -geproci, there needs to be two such cones, of degrees a, b .



Geometry gets weird in positive characteristic p ! For example, there's Fermat's Little Theorem and there's the Freshman's Dream (aka Frobenius): $(x + y)^p = x^p + y^p$. But this weirdness makes being geometric very natural!

Cones in $\mathbb{P}_{\mathbb{F}_q}^3$ of degree $a = q + 1$

Consider $Z = \mathbb{P}_{\mathbb{F}_q}^3$.

Note that $\#Z = \frac{q^4 - 1}{q - 1} = q^3 + q^2 + q + 1 = (q + 1)(q^2 + 1)$.

There is a unique degree $q + 1$ cone containing Z whose vertex is at a general point $P = (a, b, c, d) \in \mathbb{P}_k^3$, $k = \overline{\mathbb{F}_q}$. This cone meets every line through two points of $\mathbb{P}_{\mathbb{F}_q}^3$ transversely. It is given by

$$\begin{vmatrix} a & b & c & d \\ a^q & b^q & c^q & d^q \\ x & y & z & w \\ x^q & y^q & z^q & w^q \end{vmatrix} = 0$$

Is there a cone of degree $b = q^2 + 1$? There is!

Each line of $\mathbb{P}_{\mathbb{F}_q}^3$ contains $q + 1$ points. Can $\mathbb{P}_{\mathbb{F}_q}^3$ be partitioned by mutually-skew lines? Yes! Such a partition is called a **spread**, a name from combinatorics. The fibers S^1 of the Hopf fibration H map to the fibers $\mathbb{P}_{\mathbb{R}}^1$ of F , which give an example of a spread in $\mathbb{P}_{\mathbb{R}}^3$.

$$\begin{array}{ccc} S^3 & \xrightarrow{H} & S^2 \\ \downarrow A & & \downarrow = \\ \mathbb{P}_{\mathbb{R}}^3 & \xrightarrow{F} & \mathbb{P}_{\mathbb{C}}^1 \end{array}$$

For $\mathbb{P}_{\mathbb{F}_q}^3$, there are $q^2 + 1$ lines in the spread. The join of each line of the spread with P is our cone.

A Theorem

The following result gives a new method of constructing nontrivial geproci sets.

Theorem (K-)

The set of points $\mathbb{P}_{\mathbb{F}_q}^3$ is $(q + 1, q^2 + 1)$ -geproci in \mathbb{P}_k^3 , where k is an algebraically closed field containing \mathbb{F}_q .

Note when $q = 2$, we get a non-trivial $(3, 5)$ -geproci set! No nontrivial $(3, 5)$ -geproci set exists in characteristic 0 [CFFHMSS], so this is new.

Definition

A **partial spread** of $\mathbb{P}_{\mathbb{F}_q}^3$ with **deficiency** d is a set of $q^2 + 1 - d$ mutually-skew lines. A **maximal partial spread** is a partial spread of positive deficiency that is not contained in any larger partial spread.

Maximal partial spreads give a way of producing infinitely many nontrivial non-half-grids.

Theorem (K-)

The complement of a maximal partial spread of deficiency d is a non-trivial $\{q + 1, d\}$ -geproci set. Furthermore, when $d > q + 1$, the complement is a non-trivial non-half-grid.

In 1993 and 2002, Heden proved for $q \geq 7$ that there are maximal partial spreads of every deficiency d in the interval $q - 1 \leq d \leq \frac{q^2+1}{2} - 6$.

Definition

A configuration of lines \mathcal{L} in \mathbb{P}^3 is **dual-geproci** if the general projection of \mathcal{L} into a plane H is dual to a complete intersection of points in H^* .

This projection function on lines can be thought of as a function within $Gr(2, 4)$: Let $P \in \mathbb{P}^3$ and $H \in \mathbb{P}^{3*}$, then define

$$\pi_{P,H} : Gr(2, 4) \setminus \Sigma_2(\mathcal{V}_P) \rightarrow \Sigma_{1,1}(\mathcal{V}_H)$$


as

$$\pi_{P,H}(L) = \Sigma_{1,1}(\mathcal{V}_{\overline{PL}}) \cap \Sigma_{1,1}(\mathcal{V}_H),$$

where \mathcal{V}_P , $\mathcal{V}_{\overline{PL}}$, \mathcal{V}_H are any flags containing P , \overline{PL} , and H , respectively. Then a finite set $\mathcal{L} \subseteq Gr(2, 4)$ is dual-geproci if $\pi_{P,H}(\mathcal{L})$ is a complete intersection in the plane $\Sigma_{1,1}(\mathcal{V}_H)$ for a general $P \in \mathbb{P}^3$.

Complete Bipartite Graphs

So far, the only known examples of dual-geproci sets come from complete bipartite graphs in \mathbb{P}^3 . These are the equivalent of grids, because the image of \mathcal{L} is a complete intersection of two unions of lines in $\Sigma_{1,1}(\mathcal{V}_H)$.




Geproci With Infinitely-Near Points

Theorem (K-)

Let $\text{char } k = 2$. Let $Z = \{(1, 0, 0, 0) \times 2, (0, 1, 0, 0) \times 2, (0, 0, 1, 0) \times 2\}$ (where $p_i \times 2$ represents an ordinary point $p_i \in \mathbb{P}_k^3$ and a point q_i infinitely near p_i), with the infinitely-near point at each ordinary point corresponding to the tangent along the line through p_i and $(0, 0, 0, 1)$. Then Z is a $(2, 3)$ -geproci half-grid.

No $(2, 3)$ half-grid is known in characteristic 0.




Another Example

Theorem (K-)

Let

$Z = \{(1, 0, 0, 0) \times 2, (0, 1, 0, 0) \times 2, (0, 0, 1, 0) \times 2, (0, 0, 0, 1) \times 2, (1, 1, 1, 1)\}$,
with each infinitely-near point corresponding to the line containing
 $(1, 1, 1, 1)$. Then Z is $(3, 3)$ -geproci. It is a non-trivial non-half-grid.

No nontrivial $(3, 3)$ -geproci sets are known in characteristic 0.



- L. Chiantini, L. Farnik, G. Favacchio, B. Harbourne, J. Migliore, T. Szemberg, and J. Szpond. Configurations of points in projective space and their projections. [arXiv:2209.04820](https://arxiv.org/abs/2209.04820), 2022.
- O. Heden. Maximal partial spreads and the modular n -queen problem III. *Discrete Mathematics*, 243:135–150, 2002.
- O. Heden. Maximal partial spreads and the modular n -queen problem. *Discrete Mathematics*, 120:75–91, 1993.