

# Groupoids of Configurations of Lines

Jake Kettinger

Colorado State University  
Speaking at the 2025 Lefschetz Conference in Nordfjordeid

3 June 2025

## Definition

A finite set  $Z$  in  $\mathbb{P}_k^n$  is **geproci** if the projection  $\overline{Z}$  of  $Z$  from a general point  $P$  to a hyperplane  $H = \mathbb{P}_k^{n-1}$  is a complete intersection in  $H$ .

Geproci stands for **g**eneral **p**rojection is a **c**omplete **i**ntersection. The only nontrivial examples known are for  $n = 3$ . In this case a hyperplane is a plane. A reduced set of points in a plane is a complete intersection if it is the transverse intersection of two algebraic curves. For  $\#Z = ab$  ( $a \leq b$ ),  $Z$  is  $(a, b)$ -geproci if  $\overline{Z}$  is the intersection of a degree  $a$  curve and a degree  $b$  curve.

# Background: $k$ -Spreads in $\mathbb{P}_F^n$

## Definition

Given a  $k \geq 0$  and field  $F$ , a  **$k$ -spread** of  $\mathbb{P}_F^n$  is a set of mutually-skew  $k$ -planes that partition  $\mathbb{P}_F^n$ .

Spreads are known to exist for  $F =$  any finite field if and only if  $k + 1$  divides  $n + 1$ , also and for  $k + 1 = (n + 1)/2$  for  $F = \mathbb{R}$ .

Spreads are instrumental for the proof that  $\mathbb{P}_{\mathbb{F}_q}^3$  is geproci under  $\mathbb{P}_{\overline{\mathbb{F}_q}}^3$ . In this case, a spread is a partition of  $\mathbb{P}_{\mathbb{F}_q}^3$  into lines.

# The Hopf Fibration over $\mathbb{R}$

The Hopf fibration  $H : S^3 \rightarrow S^2$  can yield a spread over  $\mathbb{P}_{\mathbb{R}}^3$ .

$$\begin{array}{ccc} S^3 & \xrightarrow{H} & S^2 \\ \downarrow A & & \downarrow \cong \\ \mathbb{P}_{\mathbb{R}}^3 & \xrightarrow{F} & \mathbb{P}_{\mathbb{C}}^1 \end{array}$$

Let  $L_{a,b}$  denote the line joining the points  $(1, 0, a, b)$  and  $(0, 1, -b, a)$ , and let  $L_{\infty}$  denote the line joining  $(0, 0, 1, 0)$  and  $(0, 0, 0, 1)$ . Then  $\mathcal{S} = \{L_{a,b} : a, b \in \mathbb{R}\} \cup \{L_{\infty}\}$  is the spread yielded by the Hopf fibration. You can also use a similar method in positive characteristic to define a spread known as the **Hopf spread**.

According to Gorla, it is possible to construct a spread of  $k$ -dimensional hyperplanes inside  $\mathbb{P}_{\mathbb{F}_q}^n$  if and only if  $k + 1$  divides  $n + 1$ .

Let  $p(x) \in \mathbb{F}_q[x]$  be irreducible, monic, and degree  $k + 1$ , and let  $P$  be its companion matrix. Then one can construct a spread of spaces of the form

$$\underbrace{\text{rowsp} \left( 0 \quad \cdots \quad 0 \quad \boxed{I_{k+1}} \quad A_1 \quad \cdots \quad A_j \right)}_{\frac{n+1}{k+1}} : A_i \in \mathbb{F}_q[P].$$

When  $k = 1$  and  $n = 3$ , this construction is identical to the Hopf spread! But this also provides new examples of spreads in higher dimensions.

# Non-Hopf Spreads

## Definition

A **regulus** is a set of at least three mutually-skew lines  $\mathcal{R}$  such that there is a quadric surface  $Q$  where  $\bigcup_{R \in \mathcal{R}} R \subseteq Q$ , [like this](#).

Every regulus  $\mathcal{R}$  admits an **opposite regulus**  $\mathcal{R}^*$ .

The Hopf spread contains reguli: for example  $\mathcal{R} = \{L_{a,0} : a \in k\} \cup \{L_\infty\}$  is a regulus.

Given a spread  $\mathcal{S}$  containing a regulus  $\mathcal{R}$ , the set of lines  $(\mathcal{S} \setminus \mathcal{R}) \cup \mathcal{R}^*$  is also a spread.

It was once conjectured that every spread could be constructed by starting with the Hopf spread and replacing a succession of reguli with their opposites. This is now known to be false.

# Maximal Partial Spreads

Note that a spread over  $\mathbb{P}_{\mathbb{F}_q}^3$  comprises  $q^2 + 1$  mutually-skew lines.

## Definition

A **partial spread** of  $\mathbb{P}_{\mathbb{F}_q}^3$  with **deficiency**  $d$  is a set of  $q^2 + 1 - d$  mutually-skew lines. A **maximal partial spread** is a partial spread of positive deficiency that is not contained in any larger partial spread.

Maximal partial spreads are also instrumental to the study of geproci sets; in fact, given a maximal partial spread  $\mathcal{M}$ , the set  $\mathbb{P}_{\mathbb{F}_q}^3 \setminus (\bigcup_{L \in \mathcal{M}} L)$  is geproci.

# Projecting a Line to a Line via a Line

## Definition

Given three lines  $L_1, L_2, L_3 \subseteq \mathbb{P}_k^3$  where  $L_1 \cap L_2 = L_2 \cap L_3 = \emptyset$ , we can define the function  $\pi(L_1, L_2, L_3) : L_1 \rightarrow L_3$  as follows: take  $p \in L_1$ . Then there is a unique line  $T$  such that  $p \in T$ ,  $T \cap L_2 \neq \emptyset$ , and  $T \cap L_3 \neq \emptyset$ . Then define  $\pi(L_1, L_2, L_3)(p) = T \cap L_3$ . This is the **projection** of  $L_1$  to  $L_3$  **via**  $L_2$ .

[Here is a demonstration.](#)



## Definition

A **groupoid** is a category  $\mathcal{G}$  where every morphism is invertible.

- For any object  $G \in \mathcal{G}$ ,  $\text{Hom}_{\mathcal{G}}(G, G) = \text{Aut}_{\mathcal{G}}(G)$  is a group.  $\text{Aut}_{\mathcal{G}}(G)$  is a “group of the groupoid.”
  - Whenever  $\text{Hom}_{\mathcal{G}}(G_1, G_2) \neq \emptyset$ , then  $\text{Aut}_{\mathcal{G}}(G_1) \cong \text{Aut}_{\mathcal{G}}(G_2)$ .
- So when  $\text{Hom}_{\mathcal{G}}(G_1, G_2) \neq \emptyset$  for all  $G_1, G_2 \in \mathcal{G}$ ,  $\mathcal{G}$  induces only one group of the groupoid, up to isomorphism. Then it makes sense to say “the” group of the groupoid,  $\text{Aut}_{\mathcal{G}}$ .

## Theorem

*Let  $\mathcal{L}$  be a set of lines in  $\mathbb{P}_F^3$ . Define  $\Pi$  to be the composition-closure of the set of functions  $\{\pi(L_i, L_j, L_k) : L_i, L_j, L_k \in \mathcal{L}, L_i \cap L_j = L_j \cap L_k = \emptyset\}$ . Then  $(\mathcal{L}, \Pi)$  is a groupoid.*

In this case, any group of the groupoid is a subgroup of  $\text{Aut}(\mathbb{P}_F^1) \cong \text{PGL}(2, F)$ .

What can we say about this groupoid and its corresponding group(s)? In characteristic 0, when is it finite versus infinite? When does  $\text{Aut}_{(\mathcal{L}, \Pi)}(L)$  have finite orbits, or finitely many orbits? What is the relationship (if  $\exists$ ) between  $\text{Aut}_{(\mathcal{L}, \Pi)} \leq \text{Aut}(\mathbb{P}^1)$  and  $\text{Aut}(\mathcal{L}) \leq \text{Aut}(\mathbb{P}^3)$ ?

# Ganger's Results

In her 2024 thesis, Ganger used the technique of **transversals** to prove the following theorem:

## Theorem (Ganger Corollary 2.5)

*The group of the groupoid for the Hopf spread induced by the degree-2 field extension  $\mathbb{F}_{q^2}/\mathbb{F}_q$  over a finite field is isomorphic to the quotient  $\mathbb{F}_{q^2}^*/\mathbb{F}_q^* \cong C_{q+1}$ .*

## Definition

Given a set of lines  $\mathcal{L}$  in  $\mathbb{P}_F^3$ , a **transversal** is a line  $T$  in  $\mathbb{P}_F^3$  such that  $T \cap \bar{L} \neq \emptyset$  for all  $L \in \mathcal{L}$ .

The Hopf spread has exactly two transversals  $T_1, T_2$  for any finite field. The intersection of transversal with a line  $L \in \mathcal{S}$  is a fixed point of  $\text{Aut}_{(\mathcal{S}, \Pi)}(L)$ !

# Representing $\text{Aut}_{\mathcal{G}} \leq \text{PGL}(2, F)$

Let  $U = \overline{u_0 u_1}$ ,  $V = \overline{v_0 v_1}$ ,  $W = \overline{w_0 w_1}$  be lines in  $\mathbb{P}^3$ . Then any point on  $U$  can be written as  $au_0 + bu_1$  for  $(a, b) \in \mathbb{P}^1$  and any point on  $W$  can be written as  $cw_0 + dw_1$  for  $(c, d) \in \mathbb{P}^1$ . Then if

$$cw_0 + dw_1 = \pi(U, V, W)(au_0 + bu_1),$$

we must have  $\overline{(au_0 + bu_1)(cw_0 + dw_1)} \cap V \neq \emptyset$ .

Therefore the wedge product

$$(au_0 + bu_1) \wedge v_0 \wedge v_1 \wedge (cw_0 + dw_1) = 0.$$

We can write this as a matrix formula

$$\begin{pmatrix} a & b \end{pmatrix} \begin{pmatrix} u_0 \wedge v_0 \wedge v_1 \wedge w_0 & u_0 \wedge v_0 \wedge v_1 \wedge w_1 \\ u_1 \wedge v_0 \wedge v_1 \wedge w_0 & u_1 \wedge v_0 \wedge v_1 \wedge w_1 \end{pmatrix} \begin{pmatrix} c \\ d \end{pmatrix} = 0,$$

or rewrite this equivalently:

$$\begin{pmatrix} -u_0 \wedge v_0 \wedge v_1 \wedge w_1 & -u_1 \wedge v_0 \wedge v_1 \wedge w_1 \\ u_0 \wedge v_0 \wedge v_1 \wedge w_0 & u_1 \wedge v_0 \wedge v_1 \wedge w_0 \end{pmatrix} \begin{pmatrix} a \\ b \end{pmatrix} = \begin{pmatrix} c \\ d \end{pmatrix}.$$

## Example: The $D_4$ configuration

### Theorem

*Let  $\mathcal{L}$  be the 16 lines of the  $(12_4, 16_3)$  configuration  $D_4$  and let  $\Pi$  be the composition-closure of the projection functions. Then the group of the groupoid  $\mathcal{G} = (\mathcal{L}, \Pi)$  is  $\text{Aut}_{\mathcal{G}} \cong S_3$ .*

Argument boils down to:

- $\text{Hom}_{\mathcal{G}}(L, L') \neq \emptyset$  for  $L, L' \in \mathcal{L}$ , so  $\text{Aut}_{\mathcal{G}}$  is well-defined.
- Let  $q \in L$  be a quadruple point and  $\pi \in \text{Hom}_{\mathcal{G}}(L, L')$ . Then  $\pi(q)$  is a quadruple point. So  $\text{Aut}_{\mathcal{G}} \leq S_3$ .
- We have found automorphisms in  $\text{Aut}_{\mathcal{G}}(L)$  of orders 2 and 3, so  $\text{Aut}_{\mathcal{G}} \cong S_3$ .


In fact, one can find a subset of six lines whose group of the groupoid is  $S_3$ !

# A Helpful Labeling

Let  $\{A, B, C\}$  be a set of three letters, and consider the group  $(\mathbb{Z}/2\mathbb{Z})^2 = \langle z_1, z_2 \rangle$ . Then one can label the vertices of the  $D_4$  configuration with the elements of  $\{A, B, C\} \oplus (\mathbb{Z}/2\mathbb{Z})^2$  in such a way that there is a bijective correspondence between the lines of the  $D_4$  and triples of the form  $\{A + g, B + g', C + g'' : g + g' + g'' = 0\} \subseteq \{A, B, C\} \oplus (\mathbb{Z}/2\mathbb{Z})^2$ .

This makes the theorem on the previous slide easier to prove because you can divide the vertices into “types”  $A$ ,  $B$ , and  $C$ .

# A Helpful Labeling



- $\{A, B, C\}$
- $\{A + z_1, B + z_2, C + z_1 + z_2\}$
- $\{A + z_2, B + z_1 + z_2, C + z_1\}$
- $\{A + z_1 + z_2, B + z_1, C + z_2\}$

# Takk!

Takk så mykje!