

Groupoids of Configurations of Lines

Jake Kettinger

Colorado State University
Speaking at the AMS 2026 Spring Western Sectional Meeting

8 March 2026

Definition

A finite set Z in \mathbb{P}_k^n is **geproci** if the projection \bar{Z} of Z from a general point P to a hyperplane $H = \mathbb{P}_k^{n-1}$ is a complete intersection in H .

Geproci stands for **g**eneral **p**rojection is a **c**omplete **i**ntersection. The only nontrivial examples known are for $n = 3$. In this case a hyperplane is a plane. A reduced set of points in a plane is a complete intersection if it is the transverse intersection of two algebraic curves. For $\#Z = ab$ ($a \leq b$), Z is (a, b) -geproci if \bar{Z} is the intersection of a degree a curve and a degree b curve.

Definition

Given a $k \geq 0$ and field F , a **k -spread** of \mathbb{P}_F^n is a set of mutually-skew k -planes that partition \mathbb{P}_F^n .

Spreads are known to exist for $F =$ any finite field if and only if $k + 1$ divides $n + 1$, and also for $k + 1 = (n + 1)/2$ for $F = \mathbb{R}$.

Spreads are instrumental for the proof that $\mathbb{P}_{\mathbb{F}_q}^3$ is geproci in $\mathbb{P}_{\mathbb{F}_q}^3$ [K– 24].

In this case, a spread is a partition of $\mathbb{P}_{\mathbb{F}_q}^3$ into lines.

Gorla's Construction

It is possible to construct a spread of k -dimensional hyperplanes inside $\mathbb{P}_{\mathbb{F}_q}^n$ if and only if $k + 1$ divides $n + 1$.

Let $p(x) \in \mathbb{F}_q[x]$ be irreducible, monic, and degree $k + 1$, and let P be its companion matrix. Then one can construct a spread of spaces of the form

$$\text{row} \left(\underbrace{0 \quad \cdots \quad 0 \quad \boxed{I_{k+1}} \quad A_1 \quad \cdots \quad A_j}_{\frac{n+1}{k+1}} \right) : A_i \in \mathbb{F}_q[P].$$

Maximal Partial Spreads

Note that a spread over $\mathbb{P}_{\mathbb{F}_q}^3$ comprises $q^2 + 1$ mutually-skew lines.

Definition


A **partial spread** of $\mathbb{P}_{\mathbb{F}_q}^3$ with **deficiency** d is a set of $q^2 + 1 - d$ mutually-skew lines. A **maximal partial spread** is a partial spread of positive deficiency that is not contained in any larger partial spread.

Maximal partial spreads are also instrumental to the study of geproci sets; in fact, given a maximal partial spread \mathcal{M} , the set $\mathbb{P}_{\mathbb{F}_q}^3 \setminus (\bigcup_{L \in \mathcal{M}} L)$ is geproci [K– 24].

Projecting a Line to a Line via a Line

Definition

Given three lines $L_1, L_2, L_3 \subseteq \mathbb{P}_k^3$ where $L_1 \cap L_2 = L_2 \cap L_3 = \emptyset$, we can define the function $\pi(L_1, L_2, L_3) : L_1 \rightarrow L_3$ as follows: take $p \in L_1$. Then there is a unique line T such that $p \in T$, $T \cap L_2 \neq \emptyset$, and $T \cap L_3 \neq \emptyset$. Then define $\pi(L_1, L_2, L_3)(p) = T \cap L_3$. This is the **projection** of L_1 to L_3 **via** L_2 .



Definition

A **groupoid** is a category \mathcal{G} where every morphism is invertible.

- For any object $G \in \mathcal{G}$, $\text{Hom}_{\mathcal{G}}(G, G) = \text{Aut}_{\mathcal{G}}(G)$ is a group.

$\text{Aut}_{\mathcal{G}}(G)$ is a “group of the groupoid.”

- Whenever $\text{Hom}_{\mathcal{G}}(G_1, G_2) \neq \emptyset$, then $\text{Aut}_{\mathcal{G}}(G_1) \cong \text{Aut}_{\mathcal{G}}(G_2)$.

So when $\text{Hom}_{\mathcal{G}}(G_1, G_2) \neq \emptyset$ for all $G_1, G_2 \in \mathcal{G}$, \mathcal{G} induces only one group of the groupoid, up to isomorphism. Then it makes sense to say “the” group of the groupoid, $\text{Aut}_{\mathcal{G}}$.

Theorem

Let \mathcal{L} be a set of lines in \mathbb{P}_F^3 . Define Π to be the composition-closure of the set of functions $\{\pi(L_i, L_j, L_k) : L_i, L_j, L_k \in \mathcal{L}, L_i \cap L_j = L_j \cap L_k = \emptyset\}$. Then (\mathcal{L}, Π) is a groupoid.

In this case, any group of the groupoid is a subgroup of $\text{Aut}(\mathbb{P}_F^1) \cong \text{PGL}(2, F)$.

What can we say about this groupoid and its corresponding group(s)? In characteristic 0, when is it finite versus infinite? When does $\text{Aut}_{(\mathcal{L}, \Pi)}(L)$ have finite orbits, or finitely many orbits? What is the relationship (if \exists) between $\text{Aut}_{(\mathcal{L}, \Pi)} \leq \text{Aut}(\mathbb{P}^1)$ and $\text{Aut}(\mathcal{L}) \leq \text{Aut}(\mathbb{P}^3)$? In characteristic 0, it is known that when $\text{Aut}_{(\mathcal{L}, \Pi)}$ is finite, then orbits of points under this groupoid are geproci [CFFHMSS 25+].

Ganger's Results

In her 2024 thesis, Ganger used the technique of **transversals** to prove the following theorem:

Theorem (Ganger Corollary 2.5)

The group of the groupoid for the Hopf spread induced by the degree-2 field extension $\mathbb{F}_{q^2}/\mathbb{F}_q$ over a finite field is isomorphic to the quotient $\mathbb{F}_{q^2}^/\mathbb{F}_q^* \cong C_{q+1}$.*

Definition

Given a set of lines \mathcal{L} in \mathbb{P}_F^3 , a **transversal** is a line T in \mathbb{P}_F^3 such that $T \cap \bar{L} \neq \emptyset$ for all $L \in \mathcal{L}$.

The Hopf spread has exactly two transversals T_1, T_2 for any finite field. The intersection of transversal with a line $L \in \mathcal{S}$ is a fixed point of $\text{Aut}_{(\mathcal{S}, \Pi)}(L)$!

Representing $\text{Aut}_{\mathcal{G}} \leq \text{PGL}(2, F)$

Let $U = \overline{u_0 u_1}$, $V = \overline{v_0 v_1}$, $W = \overline{w_0 w_1}$ be lines in \mathbb{P}^3 . Then there is a unique matrix $A \in \text{PGL}(2)$ such that the following diagram commutes.


$$\begin{array}{ccc} \mathbb{P}^1 & \xrightarrow{A} & \mathbb{P}^1 \\ \downarrow \overline{u_0 u_1} & & \downarrow \overline{w_0 w_1} \\ U & \xrightarrow{\pi(U, V, W)} & W \end{array}$$

A can be taken to be $\begin{pmatrix} -u_0 \wedge v_0 \wedge v_1 \wedge w_1 & -u_1 \wedge v_0 \wedge v_1 \wedge w_1 \\ u_0 \wedge v_0 \wedge v_1 \wedge w_0 & u_1 \wedge v_0 \wedge v_1 \wedge w_0 \end{pmatrix}$.

Example: The D_4 configuration

Theorem (K-)

Let \mathcal{L} be the 16 lines of the $(12_4, 16_3)$ configuration D_4 and let Π be the composition-closure of the projection functions. Then the group of the groupoid $\mathcal{G} = (\mathcal{L}, \Pi)$ is $\text{Aut}_{\mathcal{G}} \cong S_3$.



Example: The D_4 configuration

Theorem (K-)

Let \mathcal{L} be the 16 lines of the $(12_4, 16_3)$ configuration D_4 and let Π be the composition-closure of the projection functions. Then the group of the groupoid $\mathcal{G} = (\mathcal{L}, \Pi)$ is $\text{Aut}_{\mathcal{G}} \cong S_3$.

Argument boils down to:

- $\text{Hom}_{\mathcal{G}}(L, L') \neq \emptyset$ for $L, L' \in \mathcal{L}$, so $\text{Aut}_{\mathcal{G}}$ is well-defined.
- Let $q \in L$ be a quadruple point and $\pi \in \text{Hom}_{\mathcal{G}}(L, L')$. Then $\pi(q)$ is a quadruple point. So $\text{Aut}_{\mathcal{G}} \leq S_3$.
- We have found automorphisms in $\text{Aut}_{\mathcal{G}}(L)$ of orders 2 and 3, so $\text{Aut}_{\mathcal{G}} \cong S_3$.

In fact, one can find a subset of six lines whose group of the groupoid is S_3 [K- 25+]!

Other Finite Groupoids

Using a similar analysis, we have found finite groupoids with automorphism groups isomorphic to A_4 and S_4 within the Penrose and Klein configurations [K– 25+]. These are nontrivial (5, 8)- and (6, 10)-geproci configurations, respectively.

$(1, 0, 0, 0)$	$(0, 1, 0, 0)$	$(0, 0, 1, 0)$	$(0, 0, 0, 1)$
$(0, 1, -1, 1)$	$(1, 0, -1, -1)$	$(1, -1, 0, 1)$	$(1, 1, 1, 0)$
$(0, 1, -t, t^2)$	$(1, 0, -t, -t^2)$	$(1, -t, 0, t^2)$	$(1, t, t^2, 0)$
$(0, 1, -t^2, t)$	$(1, 0, -t^2, -t)$	$(1, -t^2, 0, t)$	$(1, t^2, t, 0)$
$(0, 1, -t, 1)$	$(1, 0, -1, -t)$	$(1, -t^2, 0, t^2)$	$(1, t, 1, 0)$
$(0, 1, -t^2, t^2)$	$(1, 0, -t, -1)$	$(1, -1, 0, t)$	$(1, t^2, t^2, 0)$
$(0, 1, -1, t)$	$(1, 0, -t^2, -t^2)$	$(1, -t, 0, 1)$	$(1, 1, t, 0)$
$(0, 1, -t^2, 1)$	$(1, 0, -1, -t^2)$	$(1, -t, 0, t)$	$(1, t^2, 1, 0)$
$(0, 1, -1, t^2)$	$(1, 0, -t, -t)$	$(1, -t^2, 0, 1)$	$(1, 1, t^2, 0)$
$(0, 1, -t, t)$	$(1, 0, -t^2, -1)$	$(1, -1, 0, t^2)$	$(1, t, t, 0)$

$(0, 0, 1, 1)$	$(0, 0, 1, i)$	$(0, 0, 1, -1)$	$(0, 0, 1, -i)$
$(0, 1, 0, 1)$	$(0, 1, 0, i)$	$(0, 1, 0, -1)$	$(0, 1, 0, -i)$
$(0, 1, 1, 0)$	$(0, 1, i, 0)$	$(0, 1, -1, 0)$	$(0, 1, -i, 0)$
$(1, 0, 0, 1)$	$(1, 0, 0, i)$	$(1, 0, 0, -1)$	$(1, 0, 0, -i)$
$(1, 0, 1, 0)$	$(1, 0, i, 0)$	$(1, 0, -1, 0)$	$(1, 0, -i, 0)$
$(1, 1, 0, 0)$	$(1, i, 0, 0)$	$(1, -1, 0, 0)$	$(1, -i, 0, 0)$
$(1, 0, 0, 0)$	$(0, 1, 0, 0)$	$(0, 0, 1, 0)$	$(0, 0, 0, 1)$
$(1, 1, 1, 1)$	$(1, 1, 1, -1)$	$(1, 1, -1, 1)$	$(1, 1, -1, -1)$
$(1, -1, 1, 1)$	$(1, -1, 1, -1)$	$(1, -1, -1, 1)$	$(1, -1, -1, -1)$
$(1, 1, i, i)$	$(1, 1, i, -i)$	$(1, 1, -i, i)$	$(1, 1, -i, -i)$
$(1, -1, i, i)$	$(1, -1, i, -i)$	$(1, -1, -i, i)$	$(1, -1, -i, -i)$
$(1, i, 1, i)$	$(1, i, 1, -i)$	$(1, -i, 1, i)$	$(1, -i, 1, -i)$
$(1, i, -1, i)$	$(1, i, -1, -i)$	$(1, -i, -1, i)$	$(1, -i, -1, -i)$
$(1, i, i, 1)$	$(1, i, -i, 1)$	$(1, -i, i, 1)$	$(1, -i, -i, 1)$
$(1, i, i, -1)$	$(1, i, -i, -1)$	$(1, -i, i, -1)$	$(1, -i, -i, -1)$

A preprint is now available on the arXiv!

On to Higher Dimensions!

Giuseppe Favacchio and I are now interested in studying the groupoid of mutually-skew configurations of $n - 1$ -planes in $\mathbb{P}^{2n-1}(F)$ of the form

$$\{H(M) := \text{row} \begin{pmatrix} I_n & M \end{pmatrix}, H_\infty := \text{row} \begin{pmatrix} 0 & I_n \end{pmatrix} : M \in F^{n \times n}\}.$$

Then $\pi_{ijk} : H(M_i) \xrightarrow{H(M_j)} H(M_k)$ is represented by the matrix

$$(M_k - M_j)^{-1}(M_i - M_j) \text{ [Favacchio 25+]}.$$

We are now working on applying this construction in higher dimensions. Note $H(M) \cap H(M') = \emptyset$ iff $\det(M - M') \neq 0$.

Thank you!

Thanks for your attention! ¹

¹(Pssst, I am on the job market!)